

OUD SBIRT/Clinical Algorithm



Provide Universal SUD/OUD screening with validated tool

+ Risk factors: provide brief intervention discuss risk reduction

Screen positive SUD/OUD

OB provider to see patient, provide brief intervention to assess diagnosis, counsel risks, assess readiness for treatment (SBIRT Counseling)

Withdrawal symptoms &/or ready to start MAT

Unclear if MAT indicated, Not ready to start MAT or Outpatient MAT available

Document OUD in problem list : 099.320

Start OUD Clinical Care Checklist

Admit to hospital for Fast-Track MAT start

Initiate outpatient stabilization with Social Work support

Bill for SBIRT: < 30 min G0396 ≥ 30 min G0397

Hep C screen
Narcan Counseling
Serial Tox screen w/ consent
Neo/Peds consult
Social Work Consult
Anesthesia consult
MFM consult
Contraception counseling

Stabilize MAT and discharge to Recovery Treatment Program

Warm Handoff to Behavioral Health/ Recovery Treatment Program

Close OB follow up every 1-2 weeks (pregnancy and postpartum)

Provide standardized patient education: OUD/NAS, mom's important role in care of opioid exposed newborn (breastfeeding, rooming in, eat-sleep-console)

Inpatient Treatment Program
Intensive Outpatient Treatment
Behavioral Health Treatment Support
Peer Support Program

IL OUD Hotline
MAT/Recovery Treatment locations: 1-833-2-FINDHELP
IL Doc Assist for free Perinatal OUD Addiction Med Consult: 1-866-986-ASST (2778)